## Evolocus

Supplementary Information for

# Transgenerational epigenetic compensation and natural selection

Dmitri L. Vyssotski<sup>1,2,3</sup>

Published 17 July 2014, Evolocus 1, 19-24 (2014): http://www.evolocus.com/evolocus/v1/evolocus-01-019.pdf

Transgenerational epigenetic compensation is induced in animals by significant disturbance of parental early ontogenesis. Transgenerational epigenetic compensation consists of unblocking of one (and usually only one) previously absolutely dormant gene. This gene is selected from the whole population of dormant genes inside this organism and we can call this process "Orthoselektion" (after J.W. Harms, 1934, S. 201). Being unblocked, this gene becomes dominant immediately (can affect phenotypes of all animals where it is present in either heterozygous or homozygous state). The expression of dormant gene has two "locks": the first "lock" is removed by strong disturbance of parental early ontogenesis, the second "lock" is always present and its current state ("locked" or "unlocked") is controlled by sex-hormones and stress-hormones inside particular descendant. The expression of this gene is strictly binary (all-ornothing) and the shifting of expression episodes in time during ontogenesis can produce enormous variety of phenotypes.

Transgenerational epigenetic compensation can be induced in animals by significant disturbance of parental early ontogenesis. This disturbance can be genetic (*e.g.* presence of mutant gene in homozygous state), environmental (prenatal, neonatal or adolescent drug treatment), or their combination (exposure of a homozygous mutant to environmental stressor during early ontogenesis). The exposure of an adult animal to unusual conditions usually has either no transgenerational effect (weak stressor) or "*Tier tot*" (strong stressor).

The expression of previously dormant gene in descendants can be described by binary function (on-off; this is an easy part), but episodes of its expression can be distributed in time during ontogenesis in a very complex fashion (this is a difficult part of this process). The changes in time of expression of one previously dormant gene can produce a multitude of phenotypes.

The expression of previously dormant gene in a given descendant can be "zero" under normal conditions and "maximum" under stressor application (example: water consumption in our guinea pig). In an independent experiment, with different treatment and different previously dormant gene, its expression can be "maximum" under normal conditions and "zero" after stressor application (example: morphine analgesia during the second day of testing in the progeny of morphine-treated males). It can be "zero" in males and "maximum" in females (example:  $F_2$  generation, thyroxine experiment), or *vice versa* (example:  $F_1$  generation, morphine experiment).

All changes, distributed among multiple independent loci, known as commonly discussed epigenetic changes, like changes in DNA methylation, or histone modifications, are all secondary changes, they all are not primary heritable changes, but consequences of activation of one previously dormant gene.



Supplementary Fig. 1 | Gender-specific transgenerational epigenetic compensation in mutant mice. Heritable epigenetic compensation, developed by mutants for mutants under stressor, was introduced into all genotypes, due to breeding "at will". It has changed behaviour of mutant and wild-type females only. Natural selection was conducted by avian predators, tawny owls (*Strix aluco*), by elimination of incautious animals.

<sup>1</sup>Evolocus LLC, Tarrytown, New York, USA. <sup>2</sup>Institute of Anatomy, University of Zurich, Zurich, Switzerland. <sup>3</sup>P.K. Anokhin Institute of Normal Physiology, Russian Academy of Medical Sciences, Moscow, Russia. Correspondence should be addressed to D.L.V. (vyssotski@evolocus.com).



**Supplementary Fig. 2** |  $F_1$  and  $F_2$  descendants of drug-treated parents in the elevated plus-maze.  $F_1$  – descendants of morphine-treated (P30-P50) females, Sprague-Dawley rats<sup>25</sup>.  $F_2$  – descendants of prenatally (E8-E14) vinclozolin-treated males and females, Sprague-Dawley rats<sup>26</sup>. (**a**) Females. (**b**) Males. Both treatments entailed very similar gender-specific results in the  $F_1$  and  $F_2$  – increased caution (decreased time spent on the open arms), observed exclusively in females. Student's t-test. Mean ± SE.

The fate of mutant animals with given mutation in a natural population is determined by the efficacy of transgenerational epigenetic compensation (orthoselection and deblocking of one previously dormant gene – in homozygous mutant animals and for homozygous mutant animals) and the efficacy of natural selection (elimination of mutant and other animals from population). If orthoselection goes faster than natural selection – mutants will survive and even can replace normal animals, otherwise they will be eliminated from population completely and irreversibly.

The placement of laboratory mice into stressful semi-natural environment leads to the enhancement of viability of homozygous mutant females and, simultaneously, to the suppression of viability of wild-type females (**Supplementary Fig. 1**). The suppression was presumably achieved due to penetration of transgenerational epigenetic compensation, developed for homozygous mutants, into the animals without given mutation. This suppression is not a result of some kind of competition between animals, because heterozygous animals do not demonstrate any change in their survival during this time period. During 2 or 3 consecutive generations, the applied stressor has changed behaviour of mutant and wild-type females in the opposite direction. Further natural selection was conducted by avian predators, mainly tawny owls (*Strix aluco*).

In the **Supplementary Fig. 1** we can see population of mutant, heterozygous and wild-type mice and the role of transgenerational epigenetic compensation. This schematic picture is based on real experiment with  $Per2^{Brdm1}$  mutant mice (**Fig. 2**). Transgenerational epigenetic compensation is induced in homozygous mutant animals by harsh external conditions, presumably low temperatures and high amplitude temperature fluctuations. Some previously dormant gene has become unmasked and opened for further regulation of its expression. Its

expression was "maximum" in females and "zero" in males. This previously dormant gene remains unknown for us in this experiment.

We know that in some experiments with parental drug treatment we can see gender-specific effects in the progeny, namely increased caution in the  $F_1$  and  $F_2$  females in the elevated plus-maze (decreased time spent on open arms). No such effect was observed in males (**Supplementary Fig. 2**).

It seems that the increased caution in females can be a general consequence of orthoselection and activation of any one gene from many-many very different previously dormant genes. However here we can see that this increased caution is genotype-specific and is obvious in homozygous mutant females only, whereas the same activation of previously dormant gene leads to decreased caution in homozygous wild-type females (**Supplementary Fig. 1**).

Natural selection is conducted by tawny owls (*Strix aluco*). They are the most common aerial mouse predators in this area. Other aerial predators, like a short-eared owl (*Asio flammeus*), can take part in this process also. Aerial predators eliminate the most incautious animals from mouse population. These are the same mice that would show the longest time spent on the open arms of elevated plus-maze. And we know that so different drug treatments as adolescent (P30-P50) maternal morphine treatment and prenatal (E8-E14) vinclozolin treatment (of both parents) lead to increased caution in the  $F_1$  and  $F_2$  generation females (and only in females), observed in the elevated plus-maze as decreased open arm time (**Supplementary Fig. 2**).

Concerning experiment with vinclozolin we know that the increased caution in females is a result of increased gene expression. If we look at all genes with changed expression in different organs (**Supplementary Table 1**), we see that the number of genes with increased expression in hippocampus (in comparison with control) in increased only in females. We assume that the activation of one dormant gene leads to activation of a cascade of other genes. And most of them are usual, well-known genes, expressed in normal animals at some detectable level.

**Supplementary Table 1** | Number (*n*) of genes with changed expression

E	<u></u>		55	
12	<b>↑</b>	$\downarrow$	1	$\downarrow$
Hippocampus	1086	215	29	63
Amygdala	49	123	97	173
Hippocampus + Amygdala	1135	338	126	236
Heart	336	70	57	115
Kidney	83	67	385	340
Liver	36	63	166	100
Heart + Kidney + Liver	455	200	608	555
Ovary	211	94		
Uterus	138	141		
Prostate			276	836
Seminal vesicles			105	169
Testis			203	349
Gender-specific organs	349	235	584	1354

 $F_2$  rats, descendants of prenatally (E8-E14) vinclozolin-treated males & females (P).  $\uparrow$  - increased transcription.  $\downarrow$  - decreased transcription. Gene overlap is ignored. Brains were taken at P450 from females and at P360 from males (Skinner *et al.*, 2008)<sup>26</sup>. Other organs were taken at P120 (Skinner *et al.*, 2012)<sup>27</sup>.

#### Supplementary Table 2 | The Second Commandment

Translation		Original		
Shemot - Exodus - Chapter 20 (3-6)	3. You shall not have the gods of others in My presence.	<b>ג.</b> ל`א יִהְיֶה לְךָ אֱל ֹהִים אֲחֵרִים עַל פָּנַי:		
	<b>4</b> . You shall not make for yourself a graven image or any likeness which is in the heavens above, which is on the earth below, or which is in the water beneath the earth.	ד. ל`א תַעֲשָׂה לְךָ פֶסֶל   וְכָל תְּמוּנָה אֲשֶׁר בַּשְׁמֵים   מַמַּעַל וַאֲשֶׁר בָּאָרֶץ מִתַּחַת וַאֲשֶׁר בַּמַּיִם   מתחת לארץ:		
	5. You shall neither prostrate yourself before them nor worship them, for I, the Lord, your God, am a zealous God, Who visits the iniquity of the fathers upon the sons, upon the third and the fourth generation of those who hate Me,	، לَنَّهُ بِرَّשُرِّمَا إِنَّ إِنَّهُ مِنْ اللَّهُ بَرِيلِجِتِو ذِי אِرَ ذِرِ יְה וָה אֱל הֶיךָ אֵל קַנָּא פَ קֵד עוֹן אָבוֹת עַל בָּנִים על שַׁלְשִׁים ועל ברּעים לשַׁנָאי:		
	<b>6</b> . and [I] perform loving kindness to thousands [of generations], to those who love Me and to those who keep My commandments.	ַזּר שְׁנֵשָּׁ ם אְבָי אָבֵאָים אָאָי הָבַי וּלְשׁ`מְרֵי מִצְוֹתָי: וּ. וְעֹ־שֶׂה חֶסֶד לַאֲלָפִים לְא`הָבַי וּלְשׁ`מְרֵי מִצְוֹתָי:		
Devarim - Deuteronomy - Chapter 5 (7-10)	7. You shall not have the gods of others in My presence.	<b>ז.</b> ל`א יִהְיֶה לְךָ אֱל`הִים אֲחֵרִים עַל פָּנָי:		
	<b>8</b> . You shall not make for yourself a graven image, or any likeness which is in the heavens above, which is on the earth below, or which is in the water beneath the earth.	ח. ל`א תַעֲשֶׂה לְךָ כֶּסֶל כָּל תְּמוּנָה אֲשֶׁר בַּשָׁמֵיִם מִמַּעַל וְאֲשֶׁר בָּאָרֶץ מִתָּחַת וַאֲשֶׁר בַּמֵּים מִתַּחַת לארץ:		
	<b>9</b> . You shall not prostrate yourself before them, nor worship them, for I, the Lord your God, am a zealous God, visiting the iniquity of the fathers upon the sons, upon the third and the fourth generations of those who hate Me.	ָּאָדֶעָרָ ט. ל`א תִשְׁתַּחֲוֶה לָהֶם וְל`א תָעָבְדֵם כִּי אָנ`כִי יְה`וָה אֱל`הֶיךָ אֵל קַנָּא פּ`קֵד עֲוֹ ן אָבוֹת עַל בָּנִים ועל שׁלוּשים ועל ברעים לשׁנאי:		
	<b>10</b> . And [I] perform loving kindness to thousands [of generations] of those who love Me and to those who keep My commandments.	זע שני אָנָשָׁ בּוּבָי גַבָּצָע בְּשָּׁאָיָ י. וְעִ־שָּׁה הֶסֶד לַאֲלָפִים לְא`הָבַי וּלְשׁ`מְרֵי מִצְוֹ תָּי:		

The source of translation: http://www.chabad.org/library/bible\_cdo/aid/9881 ; http://www.chabad.org/library/bible\_cdo/aid/9969 . In colloquial Modern Hebrew, the plural לָלָדִים is generally used to refer to children (of mixed or unknown sex) while בָּלָים is generally used to refer to boys. In Christian translations, the following two simplifications are typically introduced: 1) "the sons" are replaced with "the children" (without any material basis) and 2) the continuity of generations is artificially added (without any material basis also): "visiting the iniquity of the fathers on the children to the third and fourth generation of those who hate me" (Christian Bible, English Standard Version (ESV), 2011). We do not use Christian Bible, of course. The reference to Christian Bible is provided here only due to known popularity of this deviated (inaccurate) text in the internet.

The Christian interpretation, being expressed in modern biological terms, implies that the following generations are involved: P ("parental" generation),  $F_1$ ,  $F_2$ ,  $F_3$  and  $F_4$  (four "filial" generations, *i.e.* generations of descendants). This interpretation was obtained by means of blind simplification and it is obviously wrong. We can consider at least two alternative attempts of understanding. In the first alternative attempt we can assume that this sentence can be logically divided into two semi-independent parts: 1) "visiting the iniquity of the fathers upon the sons"; 2) "visiting the iniquity [of the fathers] upon the third and the fourth generation(s) of those who hate Me". If we assume that the phrase "those who hate Me" is logically linked mainly with "generation(s)", but not with "the fathers", we can think that the following generations are involved: P,  $F_1$ ,  $F_2$  and  $F_3$  (expressed in modern terms). Thus, the fathers are "the first generation" in the text, and this generation numbering should be shifted by 1 to be compatible with modern biological terms. However there is one feature of this understanding, which makes it questionable. If we apply "those who hate Me" is applicable exclusively to "the fathers", but not to "generations". In this case all "generations" occur to be "filial" generations (generations of descendants), and in accordance with modern biological classification the following generations are present here: P,  $F_1$ ,  $F_3$  and  $F_4$ . And we remember that all above-mentioned subjects are males, *i.e.* this text describes males only.

Note that due to the increase in expression, this heritable change becomes "dominant" (detectable in phenotype of both homo- and heterozygote). Note also that the increase in expression of previously dormant gene leads to the increase of expression of many other genes, but not to their suppression. Thus, the sign of change in gene expression is not changing at all in these cascades (the increase in expression of one gene leads to the increase in expression of other genes, but not to the suppression of expression of other genes). Therefore the situation in real life is asymmetrical, in spite of the fact that from pure mechanistic viewpoint we can expect equal probability of increased and decreased expression of other genes. Note that this general increase in expression is specific for females, where we can see detectable phenotype. In males, where specific phenotype is absent, the number of genes with decreased expression is higher than the number of genes with increased expression (Supplementary Table 1). This change in gene expression in males can lead to recessive phenotype only. Remember that all above-mentioned changes in gene expression are probably secondary changes, *i.e.* they are consequences of the activation of one dormant gene.

It is interesting to note that when methylation patterns were measured by Michael K. Skinner and co-authors (2013)<sup>28</sup> at two

different, but closely related, developmental stages, namely at E13 and E16, in the  $F_2$  progeny of prenatally (E8-E14) vinclozolin-treated male and female rats, no common methylation changes were found in the primordial germ cells, with an exception of one gene, identified as *Pigb*, they all were different. It means that the demethylation of 5-methylcytosine is not a primary mechanism of dormant gene deblocking in transgenerational epigenetic compensation. The main mechanism remains unknown for us. But some of its features are already known and they are very interesting, and most of them were absolutely unexpected previously.

Observations on guinea pigs have changed interpretation of all previous experiments dramatically. In all previous experiments with transgenerational epigenetic compensation we never had non-stop indicator of any important physiological trait, available through the whole life of an experimental subject, with an exception of body weight, but it is a very generalized characteristic of an animal. We had only measurements, taken at a given time point. Only in 2012 we obtained a female guinea pig, born from mother with low water consumption and normal father, which demonstrated very high water consumption as soon as it became detectable by a direct observation. We describe this "metamorphosis" of phenotype (in this case between the mother



Supplementary Fig. 3 | Guinea pig water consumption, an episode with an abrupt increase. Experimental female was obtained from one female with naturally very low water consumption and normal male. Obtained  $F_1$  female had enormously increased water consumption, observed during her adolescence and later (up to P430, Fig. 5). Afterwards, during spring and summer, when high quality grass was available, some periods of absolutely normal water consumption have appeared stochastically. Each such period appeared and disappeared abruptly. After relatively long period of normal water consumption, during the next winter, the water consumption has jumped up again. 40-day period ( $\approx$  P615-P655) is shown here, from 2013-11-13 to 2013-12-22. All animals received the same standard food, including fresh grass that was available practically the whole year round (Tarrytown, NY).

and her daughter) as "phenotypic inversion". This animal (daughter) never showed a smooth curve of normalization of her water consumption. Instead, her water consumption was always binary, it could be either "normal" or "maximum", and the switching from one to another was obviously stress-dependent (**Fig. 5** and **Supplementary Fig. 3**).

This is an example of very sharp regulation in time. It is compatible only with a single-gene effect that can not be promoted by many independent heritable changes in many-many independent loci. The locus is only one, but its switching from "on" to "off" state in time (and *vice versa*, from "off" to "on"; very abrupt, very sharp switching) can provide unbelievable variety of phenotypes (**Supplementary Fig. 6**).

In the experiment with neonatal thyroxine treatment: previously dormant gene was activated in the  $F_1$  and  $F_2$  females and in the  $F_1$  males. However in the  $F_3$ -outcross males it was initially inactivated, but due to the stressful nature of two-way avoidance task it has become activated in the middle of 5-day training period (**Fig. 3I**). Obviously, if animal behaviour in twoway avoidance task can be strongly modulated by dynamic activation-deactivation of a single gene, any individual correlation between this behaviour and brain morphology should be absent. And, in fact, no individual correlation between hippocampal mossy fiber morphology and two-way avoidance behaviour was found in the  $F_1$ - $F_3$  generations, descendants of thyroxine-treated males, despite both traits, taken separately, had statistically significant changes (Figs. S3-S4<sup>4</sup>, S17-S19<sup>4</sup>).

In the experiment with adolescent morphine treatment: previously dormant gene was initially activated in the  $F_1$  males, but the first 10 mg/kg morphine injection plus tail-withdrawal and hot-plate testing have led to its inactivation. Therefore the analgesic effect during the second testing was normal (**Fig. 4b**). In the  $F_1$  females this previously dormant gene was inactivated (**Fig. 4a**). In the  $F_2$  females this gene was initially activated, but the first testing has led to its inactivation (**Fig. 4c**). In the  $F_2$  males all theoretically possible combinations of activationdeactivation were observed (**Fig. 4d**).

Note that these observations support the existence of two semi-independent locks of a dormant gene: the first lock is very difficult to remove and it can be removed only by very specific conditions during parental early ontogenesis; the second lock is relatively easy to remove and place back, this lock is strongly gender-dependent and stress-dependent, and the stress can be rather weak in this case. Removal and placement back of this second lock entail binary expression pattern of this previously absolutely dormant gene in time.

We suppose that rather wide variety of treatments can lead to deblocking of some previously dormant gene, whose expression was permanently switched off in a given laboratory line of animals previously, as it was supposed by J.W. Harms many years ago (1929, 1934)<sup>19,20</sup>. The choice of this gene is genotypespecific, and it is useful only for given genetic constitution (e.g.: good for given homozygous mutants, bad for wild-types). However the process of its deblocking starts as a non-specific response to stressor, as it was proposed by Hans Selye  $(1952)^{13}$ , despite he did not discuss transgenerational effects of stress, to the best of my knowledge. And it is very important that among the results of transgenerational epigenetic compensation, in the progeny, there is a batch of common non-specific changes, typical for unblocking of different dormant genes (example: increased caution in F1 and F2 females). These non-specific transgenerational changes are physiologically linked with expression of specific dormant gene. May be due to this reason, may be due to some other reasons, but their appearance in phenotype, despite they are common (non-specific), is strictly genotype-specific. I.e. above-mentioned non-specific changes can be observed only with given genotype, not with any possible genotype, where unblocked dormant gene can be introduced (introduced, for example, by means of regular breeding, like in the experiment of Serge Daan and co-authors<sup>11</sup>). Important, that

www.evolocus.com/evolocus/v1/evolocus-01-019-s.pdf



Supplementary Fig. 4 | Experimental female guinea pig (a), control females (b,c) and control male (d). Note rather relaxed posture in the control animals (b-d), and "freezing" posture in the experimental one (a), known from fear-conditioning task in rats and mice. Note also partial hair loss (a). This animal (a) does not like handling and it is relatively "wild" (it is trying to escape from human hands). Body weight: 994.4 g (a); 1370.1 g (b); 1474.6 g (c); 1533.0 g (d). Body weight is given for day 20 shown in the Supplementary Fig. 3 (2013-12-02). Photographs were taken on day 3 (2013-11-15).

non-specific phenotypic changes can be formed during different stages of ontogenesis than changes, associated with transgenerational epigenetic compensation. Common changes can be build up during early ontogenesis, when expression of dormant gene can be "On", whereas phenotypic inversion could be formed during later ontogenesis, when dormant gene can be "Off" in some animals and, thus, in these animal phenotypic inversion will not be observed at all - they will be phenotypically normal with respect to phenotypic inversion, but abnormal with respect to common (non-specific) changes - they will have these non-specific changes. Honestly, the experimental data, available today with respect to the fate of non-specific adaptive changes, are even more limited than our knowledge concerning the fate of transgenerational epigenetic compensation, typically observed in the form of phenotypic inversion. However the potential importance of non-specific adaptive changes, induced by deblocking of one previously dormant gene, in evolution is obvious. They can be used as markers by females during formation of breeding pairs, and

through this mechanism the non-random breeding can proceed in real population, where breeding "at will" is possible. Nonrandom breeding can lead, in principle, to formation of a new species. We do not have any experimental data concerning the role of transgenerational epigenetic compensation in real speciation, but, hopefully, we can use other means in order to solve this problem.

### Canalization of ontogenesis and the absence of socalled "stabilizing selection" in nature

The term "canalization of ontogenesis" and the sentence "Ontogenesis in females is better canalized than ontogenesis in males" describe situation when any disturbance of ontogenesis is handled by an animal in a way that it remains alive and able to produce offspring. This statement has nothing common with the claim that morphology or some quantitative traits remain "unchanged" or "fixed". An animal remains alive usually due to the efficacy of existing feedbacks inside the most important functional systems. These systems control the achievement of



**Supplementary Fig. 5** | Experimental female guinea pig (a) and control animals (b-d). The observed hair loss and visible morphological changes can be a result of excessive urination. These morphological changes can be a result of secondary adaptation, developed to suppress urination, but this statement can be wrong and it remains unconfirmed. Photographs made by Alexei L. Vyssotski. Green background is a chalkboard.

vital physiological results. Above-mentioned functionality has nothing common with so-called "stabilizing selection" – with elimination of extremes from the both sides of distribution curve of chosen quantitative trait (**Supplementary Fig. 7b**; see, for example: Solbrig & Solbrig, 1979, p. 146)<sup>29</sup>.

First of all, stabilizing selection, being a simplistic model, does match to nothing in nature, and, second, being modelled in laboratory, it does not lead to canalization of phenotype, so common for wild species. Even more, it was shown in 1959 by Thoday (p. 197)<sup>30</sup> in *Drosophila melanogaster* that stabilizing selection results in weak phenotype (in general weak phenotype

- animals with decreased viability), which is directly the opposite feature with respect to traits, observed in the wild nature.

"Stabilizing selection", being mathematically and logically correct, force us to assume that under stable conditions there is some "optimal" phenotype, which can be changed only due to some change in external conditions. That's wrong. First of all, as a rule, chosen indicator, convenient for measurement or observation, is linked with the most important characteristic of a given functional system in a rather correlative manner (**Supplementary Fig. 7c**). And despite this indicator is chosen



Supplementary Fig. 6 | Environmentally-sensitive timing of expression of one gene imitates multiple loci. Transgenerational epigenetic compensation makes at least one previously dormant gene available for expression. However this expression is not well optimized and it is not continuously/gradually regulated, it works as an "all or nothing gate" with two positions: 1) zero expression and 2) some technically-limited maximum. The current position of this "on-off" switch is not only gender-specific, but it is stress-sensitive, and it may be sensitive to external factors in general. Despite above-mentioned sensitivity, the answer of the gene is always 1-bit: zero expression (0) or maximum (1). The different timing of expression of one gene in four different animals (a) can completely imitate the presence of several independently modified loci, both during investigation of individual correlations (b) and during investigation of the disappearance of phenotypic traits in a raw of generations (not shown). T<sub>1</sub> & T<sub>2</sub> – ontogenetic periods of formation of traits A and B.



Supplementary Fig. 7 | Contradictory directional selection of two physiologically linked traits. Contradictory or conflicting directional selection of two physiologically linked traits (a) appears when one characteristic of an animal, let say animal size, is important for two different processes, for example for male-male competition (bigger is better) and for hiding from avian predators (the small size is easy to hide). Commonly, this situation was described as "stabilizing selection" (b), and it was supposed that it can be modelled by elimination of animals with extreme phenotypes from both sides of distribution. It was assumed that stabilizing selection is a common type of selection in nature and that it produces well canalized phenotype. Both ideas are wrong.

by a researcher as identical for two different functional systems, it is very rarely the most important dimension for both of them (Supplementary Fig. 7c). Second, the existing physiological link between different traits can be modified in evolution, it is a structure-dependent feature, and this structure can evolve. Animals are always under the pressure of mixed contradicting forces, and solution of some contradiction can be the appearance of a new species. It is not an often event, but it is always theoretically possible. All evolutionary limits, derived from the model of stabilizing selection, are false. The term "canalizing selection", as a synonym for "stabilizing selection", is also wrong, because canalization of ontogenesis is a result of directional selection, and very often a result of cyclic selection. The feature that is canalized is a set of positive results, physiological results or states, provided or achieved by phylogenetically old functional systems. Organisms that have functional systems those can achieve phylogenetically old results, despite unusual external influences, unexpected mutations and significant developmental noise, are the only ones good for further evolution. An organism can acquire in evolution something really new, only if it is able to support minimum functionality of its own old functional systems. This minimum functionality is the purpose of canalization. Morphology per se is not the object of canalization. Behaviour per se is not the object of canalization either.

Disruptive selection, as it can be organized in laboratory, is more close to real nature, than stabilizing selection, because selection of animals with opposite phenotype and their interbreeding provide load to stabilizing abilities of an organism and, as a result, the line will have improved (better stabilized) phenotype in a row of generations. This was also shown in 1959 by Thoday (pp. 196-197)<sup>30</sup>. See also the next article of Thoday & Boam (1959)<sup>31</sup>.

Concerning natural populations, Kingsolver and co-authors have shown in  $2001^{32}$ , using a bunch of published articles, that both stabilizing and disruptive selection are typically quite weak. In addition, they have shown that there is no evidence that stabilizing selection is stronger or more common than disruptive selection in nature. Taken these conclusions seriously, we can say that neither disruptive, nor stabilizing selection does exist in nature as a class. All known phenotypic results can be achieved by means of directional selection, including contradictory directional selection of two physiologically linked traits.

As we have seen previously, any randomly chosen quantitative trait, suitable for easy observation and recording, is not the thing that should be canalized. But the achievement of all positive physiological results, necessary for survival and reproduction, must be canalized.

Canalization is comprised by a set of active feedback loops. In the mechanical automates, developed by humans during previous century (XX), we typically can see one feedback loop (or, at least, one major feedback loop). In the leaving creatures, each function is supported by a set of semi-parallel feedback loops, and the inactivation one of them has relatively small negative impact on the functionality of this functional system (the impact that is usually undetectable under normal conditions). For example, the ontogenesis of females is slightly better canalized than that one of males, as it was shown by Vigen A. Geodakian in 1965<sup>33-35</sup>. It can be that some percent of semi-parallel feedback loops is disabled in males (i.e., the expression of some of their components is dependent on sex-hormones, and this expression is possible only when female sex-hormones are present and male sex-hormones are absent). We suppose, as usual, that the expression of the above-mentioned components is binary ("allor-none"): some semi-parallel feedback loops are completely disabled in males, but active in females.

We know that the transgenerational epigenetic compensation, visible at the level of phenotype as phenotypic inversion, is often present in the F<sub>2</sub> generation females, but not in the F<sub>2</sub> generation males (Fig.  $4^3$ ). It can be that this sexual dimorphism in the expression of phenotypic inversion, namely expression of some previously dormant gene in F<sub>2</sub> females, but not F<sub>2</sub> males, is the result of the same mechanism that provides better canalization of ontogenesis in females. And this mechanism is dependent on sex-hormones. For example, both previously dormant gene and some genes, involved into some semi-parallel feedback loops, can have the same regulatory sites, accessible directly or indirectly by sex-hormones. And the presence of female sexhormones allows the expression of both groups of abovementioned genes (previously dormant ones and the ones involved into feedback loops). Less simplistic view can be that both female and male sex-hormones are involved somehow. The question is, how.

We do not know the answer yet, but as a result of such functionality, females have better canalization of their ontogenesis and higher probability of expression of at least one deblocked previously dormant gene. In real life the expression of this previously dormant gene will lead to activation and expression of a lot of physiologically linked genes. We can see this, for instance, in the **Supplementary Table 1**. Note a lot of genes with increased expression in the  $F_2$  females, and only a few genes with increased expression in the  $F_2$  males; both are descendants of prenatally vinclozolin-treated males and females.

By the way, there is no contradiction or competition between "canalization" and "plasticity", because "plasticity" is not an independent entity, but it is a part of canalization process (canalization is a process, and not only a result of abovementioned process). Of course, the result of any mechanical or chemical damage *per se* is not an example of "plasticity". But the following compensation of disrupted functionality is the one.

#### **Relative fitness and selection coefficient**

Relative fitness is determined usually with respect to the most fitted phenotype, which is usually a phenotype of wild-type animal, and it is usually taken as "1" (Solbrig & Solbrig, 1979, pp. 141-142)<sup>29</sup>. Transgenerational epigenetic compensation can increase fitness of homozygous mutant phenotype so dramatically and, simultaneously, it can decrease fitness of wildtypes by a so impressive way, that selection coefficient occurs to be completely reversed (Supplementary Fig. 1). Here we can suppose that fitness is proportional to lifespan duration. Obviously, before the appearance of transgenerational epigenetic compensation, the best animals, i.e. animals with the longest lifespan, were wild-type females. After the appearance of transgenerational epigenetic compensation and after its distribution in population, the longest lifespan belongs to homozygous mutant females. And their lifespan is superior with respect to any lifespan seen before (in this experiment, of course). Thus, selection coefficient should be calculated with respect to homozygous mutant animals, starting from some time point (such time point is around generation F<sub>2</sub> in the Supplementary Fig. 1). After overwhelming distribution of transgenerational epigenetic compensation in population, the selection coefficient by chance could be even numerically the same as previously (before the appearance of epigenetic compensation), but its biological meaning will be completely

inversed. This is a result of transgenerational epigenetic compensation in evolution: it can convert homozygous mutants with previously impaired phenotype into hopeful monsters – into the animals with the best phenotype ever seen.

#### Horizontal gene transfer

Transgenerational epigenetic compensation, processing through unblocking of previously dormant genes, changes our views on horizontal gene transfer in evolution of eukaryotes. Very important role of horizontal gene transfer in evolution of prokaryotes was shown by Boris Mirkin and Eugene Koonin in 2003<sup>36,37</sup>. Transgenerational epigenetic compensation, as we see it now, does not change probability of horizontal gene transfer in higher eukaryotes directly, but it makes horizontal gene transfer less dangerous for ontogenesis and survival. If any (or practically any) transferred gene, new for a given organism, can be effectively blocked, the positive effects of horizontal gene transfer can be much higher than its negative effects.

#### Dormant genes and genetic assimilation

The deblocking of a dormant gene, being the main route of transgenerational epigenetic compensation, changes our expectations concerning the main biological function of genetic assimilation. Previously it was assumed that the genetic assimilation could be a replacement of epigenetic hereditary changes by genetic ones, and, speaking honestly, the purpose of this supposed replacement was a mystery, because it was seen as a rather complex process without clear benefits. Now we can see that the result of unblocking of one previously dormant gene is the functional appearance in the genotype of an organism of one gene with poorly determined and often unacceptable temporal pattern of expression (expression during "wrong" stages of ontogenesis and the absence of expression during "correct" developmental stages) and with poorly determined and often unacceptable dependence on sex hormones and stress hormones (expression in males instead of females, or vice versa, expression during the absence of stress instead of expression during stress, or vice versa).

The main purpose of the following genetic events, which coincide in time with the period of "genetic assimilation", is optimization of the temporal pattern of expression and optimization of sensitivity (in a very wide sense) to sex hormones and stress hormones. If the average level of expression should be changed (usually it should be decreased), it is achieved first of all by means of sinusoid-style level of expression, by means of regulation in time. And only afterwards, we can suppose, the average level of expression can be decreased in accordance with the existing requirements (but we do not have relevant experimental data now).

The question about spatial differences in expression (tissuespecific expression) seems very important, because specific expression in some structures can be extremely beneficial. Probably, the specification of the location of expression can be determined at the last steps of the discussed process, the process that can be entitled (in accordance with tradition) "genetic assimilation". Thus, the main purpose of genetic assimilation is not a replacement of transgenerational epigenetic compensation, but optimization of its results those are unexpected in many dimensions at the beginning of this process. It can be, by a lucky chance coincidence, that previously dormant gene will be absolutely ready for normal usage immediately, without any change in the functionally-linked transcription factors and other neighbouring elements. However more realistic expectation is that at least some changes will be highly desirable.

#### The priority of "dormant genes"

We assume that the priority of dormant genes belongs to J.W. Harms, despite he did not use the term "dormant gene", but, instead of this, he was speaking about "active" and "passive" genes. However he discussed "active" and "passive" genes at the levels of both genotype and phenotype, he discussed an activation of a "passive" gene by external environmental factors, and above-mentioned activation was useful for a given organism. It is important that above-mentioned activation was supposed to be heritable. He discussed an activation of a "main", the most critical, gene among a set of passive genes, and he has introduced the term "orthoselection" to describe activation of one gene among lots of irrelevant passive genes. He supposed that activated gene can control a group of other genes with their associated phenotypic traits. And he has assumed that abovementioned currently passive gene was used during previous evolutionary history of this species, together with abovementioned cascade of genes, functionally linked with each other, that is why "orthoselection" [in German: "Orthoselektion"].

There are at least two other opinions about priority of "dormant genes". Eva Jablonka and Marion J. Lamb (2014)<sup>17</sup> assume (p. 14<sup>17</sup>) that the basis of the idea of dormant genes was expressed by Darwin in his theory of "pangenesis", included in his book "*The Variation of Animals and Plants under Domestication*" (1868)<sup>38</sup>, where he described hypothetic hereditary particles – "gemmules" and stated that some gemmules can be dormant, and nutrition or climate could reawaken dormant gemmules. If we assume that a "gemmule" is a sufficient prototype of "gene", we can agree with abovementioned claim. However it seems that the gap between "gene" and "gemmule" in the ideation space is too wide for such generalization.

Dmitry K. Belyaev and co-authors  $(1981)^{21,22}$  have provided references with respect to dormant genes to publications of Muller  $(1932)^{39}$  and Zuckerkandl & Pauling  $(1962)^{40}$ . I have read the article of Muller (1932) from the beginning to the end and I have found nothing that can be associated with "dormant genes". There is a statement that there are mutations which lead to activation of some genes, *i.e.* lead to the appearance of some previously absent functionality, but that is all (see section "*Neomorphic mutations*", pp. 245-249<sup>39</sup>). Article of Zuckerkandl & Pauling (1962)<sup>40</sup> was published too late, really too late in view of publications of J.W. Harms (1929, 1934)<sup>19,20</sup>, and just due to this reason they can not pretend to any priority.

Despite the article<sup>19</sup> and the book<sup>20</sup> of J.W. Harms were unknown to D.K. Belyaev, the role of Dmitry Belyaev *per se* in the story of dormant genes should not be neglected. It was Dmitry K. Belyaev who has written the following.

1. "Once entered into an active state, the gene subsequently behaves like a discrete Mendelian unit. The idea of functional activation-inactivation of genes has permeated through the literature. It has been shared by Muller<sup>39</sup> and explicitly stated as

the dormant gene hypothesis by Zuckerkandl and Pauling<sup>40</sup>. There was no experimental substantiation of this idea.

As suggested by *star* studies, once brought out of dormancy, a gene would become functionally active and relatively stably inherited in the new state; sometimes it would become inactive, thereby distorting classical segregation patterns. The phenomenon of gene activation-inactivation cannot be explained only by the influence of the genotypic environment for the reason that within the same genotypic environment (the same organism) one of the homologous genes can be active and the other inactive" (p. 273<sup>22</sup>).

2. "The explanation offered relies on the supposition that inactive ("dormant") genes may be activated and reverted toward an inactive state. A similar reliance on gene activation-inactivation has been implied by many investigators, including Muller<sup>36</sup>. Since then, Zuckerkandl and Pauling<sup>40</sup> formulated it in their "dormant gene" hypothesis, and Belyaev<sup>41</sup> extended it to the process of domestication.

It is worth emphasizing that "dormant" genes are activated by evolutionary stimuli eventually resulting in profound metabolic changes in the cell. With regard to mice, a strong stimulus has been selection in the course of domestication" (p.  $112^{21}$ ).

3. "One can imagine that under the influence of an altered hormonal equilibrium a number of "dormant" genes could be expressed, resulting in a high frequency of appearance of a whole complex of morphological characters (position of the tail, ears, etc.) mentioned above" (p.  $306^{41}$ ).

Emile Zuckerkandl and Linus Pauling have described dormant genes in 1962<sup>40</sup> in the article "Molecular disease, evolution, and genic heterogeneity", seemingly without prior knowledge about publications of J.W. Harms (1929<sup>19</sup>, 1934<sup>20</sup>). Their description is provided in the section "D" of above-mentioned article. The section "D" is entitled "The destiny of duplicate genes and the function of genic multiplicity" (pp. 206-213<sup>40</sup>). Pages 206-210<sup>40</sup> are reprinted below as they were published in 1962 (only the reference numbers are added by us in order to be compatible with our reference style).

P. 206:

"D. The Destiny of Duplicate Genes and the Function of Genic Multiplicity

If gene duplication is one of the means of increasing the output of a given protein, one may distinguish two phases in this respect. Up to an optimum number of duplications, the duplicate gene will be selectively retained with a structure identical to and a position rather near to that of the mother gene. Beyond this point, duplicate genes will be progressively more strongly selected against. During this latter phase they will be in part eliminated with their carriers, and in part subjected to progressive change. When they are preserved and changed, their destiny may be of three types. They might evolve new useful functional properties. In this case they will be retained as active genes, and to the extent to which polypeptide output depends on gene duplication their own duplicates will be kept unchanged by natural selection. Secondly, functionless or unfavourable duplicates will not maintain duplicates to their own likeness and may be themselves be translocated to other chromosome parts and be reduced to minor-component genes by a position effect. Some such minor-component genes, more or less profoundly

changed in the meantime, may be selected for later in evolution and be changed"

P. 207:

"into major-component genes. Thirdly, the activity of the duplicate may be reduced to zero.

This elimination of gene activity may again take three forms. The changed structural gene may be bodily eliminated through the loss of the part of the chromosome that carries it; or it may be modified to such an extent that its products, although significant in amount, are no longer recognizable in terms of the original protein; or it may be preserved in a modified state, but totally or subtotally deprived of the power of expression.

The existence of such "dormant genes", although difficult to verify, is a plausible inference from two types of observations: firstly, that within a given tissue, say the hematopoietic tissue, major and minor structurally distinct components of a given type of polypeptide chain are found. Since at a given time the relative quantities of hemoglobin chains vary between 100 and 1 %, other genetically distinct minor components may be present in such small amounts that they are practically undetectable. Secondly, there are numerous examples of proteins that are produced exclusively in one type of tissue, and of which no trace is found with presently available analytical means in other tissues. The nonproduction of hemoglobin in muscle sells and of myoglobin in reticulocytes is one example. This example shows that some among even relatively closely related structural genes may, within a given tissue, be the ones strongly expressed, the others unexpressed. We must assume that all the structural genes have during embryological development been communicated to all cell lineages. It is therefore quite likely that there exist in every organism numerous structural genes that do not find in any of the existing tissues conditions favorable to their expression and thus remain permanently dormant.

Furthermore the relative structural similarity of minor hemoglobin components to one of the major components affords yet another argument in favor of the existence of dormant genes. Indeed, as we have seen, the human  $\delta$ - and  $\beta$ -chains are quite similar. Likewise structurally distinct components of orangutan hemoglobin have been found to be quite similar, and the same holds for pig hemoglobin components (E. Zuckerkandl, R. T. Jones, Y. Nishiwaki and L. Pauling, 1959-1961, unpublished). If duplicate genes remained usually expressed, one would expect to find a series of minor-component chains differing from all the other chains as much as the human  $\beta$ -chain. This is, however, not"

P. 208:

"the case. One is led to think that, in the long run, duplicate minor-component genes most often cease to be expressed. There is no apparent reason why one should assume that they have most times been bodily eliminated. Other possibilities are that they have been implied in a further translocation, that a mutation or transposition of a controlling element (repressor) has occurred, or that the structurally modified mutated gene possesses specificity characteristics that fail to comply with the specificity requirements for polypeptide production under the conditions prevailing in the cell.

Dormant genes of course are conceived as dormant only as far as their expression, and not as far as their mutability goes. Mutations in dormant genes and in minor-component genes will never be lethal, unless the latter have some distinct specific function, which would then lead us to consider them as "major components" of another protein type. Minor-component genes and, mainly, dormant genes may thus furnish an important and perhaps the principal part of the genic raw material for macroevolutionary experiments of nature. A new translocation, or the transposition of a controlling element such as those described by McClintock (1956)<sup>42</sup>, or some other genetic modification may reactivate the dormant gene after a very long period of time during which mutations have changed it enough so that it now controls the production of a new kind of protein. In this fashion new enzymes, new functions can arise without the corresponding loss of old enzymes, old functions. We have recalled earlier the importance to evolution of the loss of functions through the mutations of active genes. But it is evident that evolution, while it makes the best of such losses and of molecular disease, could not be based on them along. There are a great many more different functions to be carried out by a great many more different types of enzymes than we allowed to suppose can have existed in early evolutionary times. Primordial living matter must have been limited to a few simple functions. Therefore the notion of evolution by gain is a necessary complement to the notion of evolution by loss.

Horowitz (1945)<sup>43</sup> made a lasting contribution to our thinking about evolutionary gain at the enzymatic level. He described how new reaction chains might arise in certain circumstances through the chance combination of the necessary genes and furthermore proposed a general mechanism for the stepwise building up of complex enzyme-systems, presenting us with a plausible scheme of macro-evolution at the molecular level. Obviously, as enzyme systems"

P. 209:

"become more complex, more different enzyme molecules are needed. There are reasons to think that the same molecules cannot usually be expected to carry out several different enzymatic functions. Therefore, new genes are needed for the building up of new functions. This is where the concept of the mutational reactivation of dormant genes complements Horowitz's picture. Minor-component genes are not to be excluded from a similar role, but may usually not yet be different enough from their parental genes to be fit for carrying out novel functions. Most minor-component genes, as stated above, are liable to be eventually turned into dormant genes because natural selection will not prevent their transfer to synthetically inactive chromosome regions, nor their coming otherwise under the influence of a repressor gene, nor structural changes that might place the genes outside the range of specificity requirements of the available macromolecules that collaborate with the structural gene in protein synthesis. As dormant genes become reactivated after periods of cryptic existence corresponding perhaps to geological ages, they may produce potential enzymes that do not disrupt existing chains of reactions, but are able to add new processes to the old ones, perhaps in the ways described by Horowitz. One of the possible mechanisms of the reactivation of dormant genes is the reactivation of the chromosomal region where it is located through a change in intracellular environment. The hope of demonstrating the existence of dormant genes rests on this possibility. Such a change in intracellular environment could result from the adaptation of the organism to changes in external milieu. In adaptational changes, during an initial phase, gain and loss mutations may be balanced, or loss mutations only

may occur, so that the total complexity of the organism either remains constant or tends to decrease. In the process, however, the environment of the chromosomes may be altered in some tissue and, on account of this alteration, genes be activated in some regions of the chromosomes, inactivated in others. Inactivation of this kind will be mostly lethal and genomes with corresponding inactivation-resistant chromosome regions will be selected for. The newly activated genes on the other hand will now be available to respond to further adaptational needs and will furnish a series of gain mutations without corresponding losses. Parts of this concept are supported by observation. Changes in conformation of the genic DNA molecules are presumably related to changes in the activity of the genes, and Schmitt (1956)<sup>44</sup> has shown that the state of chromosomal DNA" P. 210:

"depends on the chemical environment of the chromosomes. Furthermore, genes in mice that display the same activity characteristics during the individual's life time have been found to be closely linked on the same chromosome (Paigen and Noell, 1961)<sup>45</sup>.

Through the reactivation of dormant genes by a modification of the intracellular environment an initial adaptational stress of great magnitude would appear instrumental in producing a rise in complexity of the organism. Thus would be solved the old paradox expressed in this question: why should organisms get to be more complex, since simpler organisms are evidently adapted as well to the environment? Once more Biology will show that it can do without any "élan vital" or "entelechy".

The present concept leads one to predict that the fastest evolution toward more highly organized forms would take place after the occurrence of major environmental stress. Evolutionary history bears out this expectation. For a long time paleontologists have noted that the initial phases in the development of new types of forms has an "explosive" character (consult, for instance, Rensch, 1954)<sup>46</sup>. According to the present theory, we assume that at the onset of such an explosive evolutionary phase a change in intracellular environment has brought about the reactivation of a number of previously dormant genes. Rensch believes that the phenomenon of explosive evolutionary phases is adequately accounted for by increased natural selection accompanying the conquest of new biotopes. He thus considers as instrumental a change in external environment only. This however does not explain the trend toward increased complexity of the forms.

In the sense that has been laid out here a marked change in environmental conditions may lead to what a layman would call a shake-up of the genome, and this may be the part of reality behind the poorly documented and falsely interpreted observations of Soviet anti-geneticist A. Lysenko.

To sum up, mutations of active genes controlling "major" protein components suffice to explain how an organism can adapt to changes in its environment. Mutations of minorcomponent genes and dormant genes, however, seem to be able to furnish the organism with the genic raw materials that eventually allow it not only to adapt to a new environment but also, in the process, to become more highly organized. Minorcomponent genes and dormant genes may thus prepare the major steps in evolution."

The concept of dormant genes, provided by Emile Zuckerkandl and Linus Pauling, is expressed in more modern terms than the article and book, written in 1929 and 1934, respectively, by Wilhelm Jürgen Heinrich Harms. However some major and minor remarks, given form the view from the 21<sup>st</sup> century, might be useful. "Dormant gene" is a catchy term. However the term "genetic locus", proposed instead of the term "gene" by James A. Shapiro in 2011 (p. 3047), provides more precise description of real situation in the field of transgenerational effects. The activation of a dormant genetic locus does not lead to the expression of this gene at some constant level. Instead, it will be expressed only if necessary combination of stress-hormones and sex-hormones will be present in a given descendant at a given time. During some generation  $(e.g., F_2)$  and in some gender (e.g., males), the gene, being brought out of absolute dormancy, and being potentially dominant, will be completely unexpressed during the whole life cycle, due to the lack of combination of stress-hormones and sex-hormones, necessary for its expression. Thus, dormant genetic locus, being brought out of dormancy, becomes open for further regulation of its expression, but not for unconditional presence of its product in the phenotype. As we have seen in the experiments with morphine and thyroxine, together with observations on guinea pigs, the expression of previously dormant genetic locus is obviously gender-dependent, significantly stress-dependent, and is sharply regulated in time during lifespan of a single organism through the effects of short periods of stress.

"Dormant genetic locus" can be not only the whole dormant gene together with all its *cis* and *trans* regulatory sites, but it can be only a part of its regulatory sites. Thus, dormant genetic locus can contain a part of regulatory sites of a given gene. Even in this case the effect of this genetic locus is strictly binary: it can be dormant and coexisting with normal level of expression of a given gene, or, alternatively, it can be brought out of dormancy and (an additional condition) in the presence of necessary stresshormones and sex-hormones it can entail very high, enormously high, level of expression of this gene.

The case, when previously dormant genetic locus will entail decreased expression, is not interesting for us and for evolution, because it will be a recessive trait, undetectable in the heterozygous organism. The asymmetry between activation and repression is very important here. If we will speak only about "regulation of expression" in general terms, we will miss the most important features of the process.

The role of "junk DNA", including the role of potentially regulatory sites in it, was discussed in the article of Emile Zuckerkandl and Giacomo Cavalli "Combinatorial epigenetics, "junk DNA", and the evolution of complex organisms" (2007)<sup>48</sup> from similar standpoint. Of course, we can discuss "the transcriptional rate of individual genes" in general. However we should remember that the transcriptional rate has minor importance, unless it is increased by a factor of four, at least. And then, when it is increased, it can be regulated only in "all-or-none" fashion and its ontogenetic windows of "on" *vs.* "off" states is the most important. Then, logically, tissue-specific expression must be optimized. We suppose that time-specific expression and stress-dependent expression will be optimized first of all.

We know that the primary removal of dormancy is a sitespecific effect, the effect which is associated with a given locus of a given chromosome. Experiments of Dmitry K. Belyaev have shown that the same gene can be dormant in one chromosome and, simultaneously, it can be expressed in homologous chromosome of the same organism. Thus, extra-chromosomal factors and intracellular conditions can not be solely responsible for above-mentioned effect.

We have not material basis to claim that the heritable unblocking of dormant genetic locus is solely epigenetic effect in modern sense, *i.e.* that it does not formed by a change of DNA sequence in a given locus. Alternatively, taking into account known variety of natural genetic engineering systems, summarized by James A. Shapiro<sup>47</sup>, it can be equally supposed that site-specific reversible genetic modification of DNA can take place with a help of specific proteins. The involvement of proteins in general in this process was shown by the experiments with Hsp90 (for a short summary, see pp. 262-263<sup>17</sup>).

Interesting that even if it will be shown that site-specific DNA sequence modification is involved as a primary factor, we still will be able to use the term "epigenetic", but in this case only in the Waddington's wide sense (not in the modern sense, given, for example, in the abstract of our current article).

Previously, the hypothesis of demethylation of 5'methylcytosine in specific DNA regions was considered as a parsimonious explanation of transgenerational effects in mammals. However it was rejected by recent experiments with vinclozolin, conducted by Michael K. Skinner and co-authors in  $2013^{28}$ . These experiments have shown that methylation patterns in the germ cells of F<sub>2</sub> generation are very different at E13 and E16 (they are so different that the demethylation process can not be considered as a primary activator of dormant genetic loci in evolution anymore).

Zuckerkandl and Pauling discussed "mutational reactivation of dormant genes" (p.  $209^{40}$ ) and "reactivation of dormant genes by a modification of the intracellular environment" (p.  $210^{40}$ ). J.W. Harms described reactivation of passive genes as an adaptive response to a new environment. This transgenerational adaptive genes and further hereditary transmission of the activated state of a chosen gene to the next generation (p.  $201^{20}$ ). Harms did not discuss the situation when the dormancy of a passive gene can be completely removed, but the gene still will not be expressed in the progeny due to the lack of some gender-related or stress-related activation factors. Zuckerkandl and Pauling did not discuss this possibility as well.

We assume that the priority of "dormant genes" is currently held by J.W. Harms, but we can change our opinion, if some other publication with early priority date will be found. It is interesting to note that Richard Goldschmidt  $(1940)^{49}$  described theoretical views of J.W. Harms the following way: "Harms explained the whole set of adaptive evolutionary changes on a purely Lamarckian basis; we refrain from discussing his arguments, which may be used as a fine example of the fallacies of that doctrine" (p. 276<sup>49</sup>).

Trofim D. Lysenko, mistakenly called by Zuckerkandl and Pauling as "A. Lysenko", discussed "unstable heredity" [in Russian: "расшатанная наследственность"] in his own publications a lot (see the term "destabilized heredity" in the Subject Index of his book "Agrobiology", published in English in 1954<sup>18</sup>). Concerning the "poorly documented observations", one can mention that they were rather "poorly accepted by Soviet Morganists" than "poorly documented". "Soviet Morganists", of course, had no relation to Thomas Hunt Morgan, the author of many articles, including popular "The theory of the gene"  $(1917)^{50}$  and "Chromosomes and heredity"  $(1910)^{51}$ . Whether "Soviet Morganists", existed in 1962, had read Morgan's articles? It is a difficult question. We know how they have read Darwin's books: they widely exploited the term "Darwinism" as the opposite to "Lamarckism", without any knowledge that Darwin himself supported the idea of "the inheritance of acquired characters"<sup>38</sup>. Contrary to many "Soviet Morganists", Trofim D. Lysenko discussed Darwinism not only as "natural selection", but as real teaching of Charles Darwin<sup>38</sup>. For an objective and compact description of Darwinism, see section "Darwin's Darwinism" (pp. 10-16<sup>17</sup>) in the book of Eva Jablonka and Marion J. Lamb "Evolution in Four Dimensions: Genetic, Epigenetic, Behavioral, and Symbolic Variation in the *History of Life*", revised edition (2014)<sup>17</sup>. Views of T.D. Lysenko can be found in a compact form in his publication "The Situation in Biological Science" (1948), see pp. 515-554<sup>18</sup>, one can look at p. 537<sup>18</sup> specifically for "destabilized heredity".

Concerning "entelechy" the situation should not be simplified too much. Just because if we have two species of animals and one of them believes into "entelechy", and another one does not believe in such non-material entities, the first one will evolve faster and in more efficient way. Especially if species are faced with the task that can not be solved during lifespan of one generation and the efforts of several consecutive generations are required. Without believe into "entelechy" an animal will not have any motivation even to try to find a solution of the task that is obviously too difficult for her. However, if she believes that her attempts, described in details, will be used as a basis for further attempts of further generations, even if all her abovementioned attempts will never be included into the "gold fund". and will be just all wrong, all in wrong direction, all completely rejected, but showing that particular field of possibilities is carefully and honestly investigated with final negative result, she may start more enthusiastically, just because our knowledge of honestly excluded possibilities may help us to find a positive solution. By the way, such animal as "rat" (as well as "mouse" and "guinea pig") is always "she" in Russian, even if we are talking about male rat (or male mouse, or male guinea pig).

Further details can be found in the books of Henri Bergson  $(1907)^{52}$ , Genrich Altshuller (1973, *e.g.* pp. 25, 48, 52)<sup>53</sup> and Ludwik Fleck  $(1935)^{54}$ .

Sometimes we can speak about the necessity of a "new synthesis"<sup>16</sup>. But may be "natural selection" and Darwin's Darwinism are already upside down? Transgenerational epigenetic compensation and dormant genetic loci have placed "natural selection" exactly into the above-mentioned position. If some population has a mutation, slightly deleterious in homozygous subjects and practically neutral in heterozygous ones, then if this population will be placed in a hard conditions, the conditions, difficult for survival, then, in accordance with the laws of natural selection, this mutation will be wiped away from the population together with the bearing it animals.

However, in real life, the hard environmental conditions will bring out of dormancy at least one genetic locus, only in the above-mentioned homozygous mutants, and we will have in population not only one previously known recessive mutation, but, in addition, one unexpected dominant mutation. Original mutant gene is typically recessive, whereas the deblocked genetic locus always bears dominant trait. Of course, original mutant gene and deblocked genetic locus are not necessarily homologous (as it was discussed by Zuckerkandl and Pauling), but they are participants of the same functional system. We use the term "functional system" exactly as it was proposed by Peter K. Anokhin (1974)<sup>55</sup>, but not in the sense of "systems biology".

The old mutant gene (*e.g.*, Per2 in the following example) and the unblocked previously unknown dormant gene are not necessarily homologous. However if they are homologous due to some unexpected lucky coincidence, it will simplify further experimental investigation enormously.

The deblocked genetic locus may change the direction of natural selection towards homozygous mutants, bearing original mutation, exactly as it is shown in the Fig. 2 and Supplementary Fig. 1. These figures show the experimental data with  $Per2^{BrdmI}$  mutant mice under semi-natural outdoor conditions, obtained in the experiment of Serge Daan and coauthors, first described in 2011<sup>11</sup>. In the most interesting case this situation can develop into a speciation (formation of a new species on the basis of homozygous mutants having independent deblocked genetic locus). Such homozygous mutants can be called "hopeful monsters", as it was proposed by Richard Goldschmidt many years ago<sup>49,56</sup>. Above-mentioned new species will be reproductively separated form the old one. The old species consists of wild-type animals without any change in the above-mentioned dormant genetic locus. This possibility was described in the section "The main modus of macroevolution" in our article "Transgenerational epigenetic compensation and sexual dimorphism"<sup>3</sup>.

#### Supplementary References (references 1-24 are given in our main paper)

- Byrnes, E.M. Transgenerational consequences of adolescent morphine exposure in female rats: effects on anxiety-like behaviors and morphine sensitization in adult offspring. *Psychopharmacol.* 182, 537-544 (2005).
- Skinner, M.K., Anway, M.D., Savenkova, M.I., Gore, A.C. & Crews, D. Transgenerational epigenetic programming of the brain transcriptome and anxiety behavior. *PLoS ONE* 3(11), e3745 (2008).
- Skinner, M.K., Manikkam, M., Haque, Md. M., Zhang, B. & Savenkova, M.I. Epigenetic transgenerational inheritance of somatic transcriptomes and epigenetic control regions. *Genome Biol.* **13**, R91 (2012).
- Skinner, M.K., Guerrero-Bosagna, C., Haque, M., Nillson, E., Bhandari, R. & McCarrey, J.R. Environmentally induced transgenerational epigenetic reprogramming of primordial germ cells and the subsequent germ line. *PLoS ONE* 8(7), e66318 (2013).
- Solbrig, O.T. & Solbrig, D.J. Introduction to Population Biology and Evolution (Addison-Wesley, Reading, MA, 1979).
- Thoday, J.M. Effects of disruptive selection: 1. Genetic flexibility. *Heredity* 13, 187-203 (1959).
- Thoday, J.M. & Boam, T.B. Effects of disruptive selection: 2. Polymorphism and divergence without isolation. *Heredity* 13, 205-218 (1959).

- Kingsolver, J.G., Hoekstra, H.E., Hoekstra, J.M., Berrigan, D., Vignieri, S.N., Hill, C.E., Hoang, A., Gibert, P. & Beerli, P. The strength of phenotypic selection in natural populations. *Am. Nat.* **157**(3), 245-261 (2001).
- Geodakian, V.A. The role of sexes in transmission and transformation of genetic information. *Problems of Information Transmission* 1, 105-112 (1965). [Publ. in Russian: Геодакян В.А. Роль полов в передаче и преобразовании генетической информации. *Пробл. Передачи Информ.*, Том 1, Вып. 1, стр. 105-112 (1965)]. www.evolocus.com/publications/geodakian1965.pdf
- Geodakian, V.A. Why two sexes? *Nauka i Zhizn (Science and Life)* No 3, 99-105 (1966). Translated from Russian by Boris Lubachevsky in 1988. www.evolocus.com/publications/geodakian1966.pdf
- Geodakian, V.A. Two Sexes. Why? The Evolutionary Theory of Sex (Ed. by Sergey Geodakyan, Wilmington, 2012). www.evolocus.com/textbooks/geodakian2012.pdf
- Mirkin, B. & Koonin, E. A top-down method for building genome classification trees with linear binary hierarchies. In *Bioconsensus*, Janowitz, M., Lapointe, J.-F., McMorris, F., Mirkin, B. & Roberts, F., Eds. (DIMACS Series, Vol. 61, AMS, Providence, 2003), pp. 97-112.
- Mirkin, B.G., Fenner, T.I., Galperin, M.Y. & Koonin, E.V. Algorithms for computing parsimonious evolutionary scenarios for genome evolution, the last universal common ancestor and dominance of horizontal gene transfer in the evolution of prokaryotes. BMC Evol. Biol. 3, 2 (2003).
- Darwin, C. The Variation of Animals and Plants under Domestication. 2 vols. 2<sup>nd</sup> ed. (Johns Hopkins Univ. Press, Baltimore, MD, 1998).
- Muller, H.J. Further studies on the nature and causes of gene mutations. Proc. Int. Congr. Genet. 6th. 1, 213-255 (1932).
- Zuckerkandl, E. & Pauling, L. Molecular disease, evolution, and genic heterogeneity. In *Horizons in Biochemistry*, Pullman, M. & Kasha, B., Eds. (Academic Press, New York, 1962), pp. 189-225.
- Belyaev, D.K. Destabilizing selection as a factor in domestication. J. Hered. 70, 301-308 (1979).
- McClintock, B. Controlling elements and the gene. Cold Spring Harb. Symp. Quant. Biol. 21, 197-216 (1956).
- Horowitz, N.H. On the evolution of biochemical syntheses. Proc. Natl. Acad. Sci. USA 31, 153-157 (1945).
- Schmitt, F.O. Patterns of interaction of biological macromolecules in relation to cell function. Proc. Natl. Acad. Sci. USA 42, 806-810 (1956).
- Paigen, K. & Noell, W.K. Two linked genes showing a similar timing of expression in mice. *Nature* **190**, 148-150 (1961).
- Rensch, B. Neuere Probleme der Abstammungslehre: Die Transspezifische Evolution, 2<sup>nd</sup> ed (Ferdinand Enke Verlag, Stuttgart, 1954).
- Shapiro, J.A. Evolution: A View from the 21<sup>st</sup> Century (FT Press Science, Upper Saddle River, NJ, 2011).
- Zuckerkandl, E. & Cavalli, G. Combinatorial epigenetics, "junk DNA", and the evolution of complex organisms. *Gene* **390**, 232-242 (2007).
- 49. Goldschmidt, R. The Material Basis of Evolution (Pageant Books, New Jersey, 1960).
- 50. Morgan, T.H. The theory of the gene. Am. Nat. 51, 513-544 (1917).
- 51. Morgan, T.H. Chromosomes and heredity. Am. Nat. 44, 449-496 (1910).
- 52. Bergson, H. Creative Evolution (Dover Publ., Mineola, New York, 1998).
- Altshuller, G.S. The Innovation Algorithm (Technical Innovation Center, Worcester, MA, 2007). Translated by L. Shulyak & S. Rodman from the Russian Second Edition, 1973; original author's title: "The Algorithm of an Invention" [Алгоритм изобретения].
- Fleck, L. Genesis and Development of a Scientific Fact. Translated by F. Bradley & T. J. Trenn (University of Chicago Press, Chicago, 1979).
- Anokhin, P.K. Biology and Neurophysiology of the Conditioned Reflex and its Role in Adaptive Behavior (Pergamon Press, New York, 1974).
- 56. Goldschmidt, R. Some aspects of evolution. Science 78, 539-547 (1933).